



Move it Monday

Physical activity improves your overall strength, balance and wellbeing – helping you stay independent.

Get moving by...

- asking your health professional for advice before you get started
- starting small and building up over time
- doing what you can, without overstretching yourself.

Why not try...


- taking a walk
- playing with grandchildren
- chair exercises supportive and easy on the joints
- hydrotherapy low-impact, full-body and good for the heart
- tai chi for balance, mindfulness and relaxation
- gentle stretching or yoga for balance and flexibility
- strength-building programs for your muscles.

Remember – any physical activity, even while sitting in a chair, is better than none!



Aim for 30 minutes of physical activity a day – this can include any form of movement, be it socialising, walking or recreational activities.

The benefits











Teach back Tuesday

Let's get educated on falls and falls prevention!

In WA, it matters because

- someone is admitted to hospital due to a fall-related injury every 15 minutes
- almost one in 5 hospitalisations of people aged 85 years or over are due to falls
- over half of all fall-related hospitalisations are due to falls in homes or aged care homes.


Reduce your risk

- talk to your health professional
- complete the Stay On Your Feet[®] checklist to know your risk of falling
- · move your body and keep active
- improve your physical and mental health
- remove trip hazards from your room
- wear appropriate shoes
- · use assistive technology
- teach someone else about falls prevention.



Knowing what to do if you fall is important. Talk to your health professional and make a plan that is best for you.

If you fall









stayonyourfeet.com.au

For more information or enquiries, contact healthpolicy@health.wa.gov.au





No Falls November Wobbly Wednesday

Postural (orthostatic) hypotension is when your blood pressure drops after you change position, such as when you move from lying to sitting, or sitting to standing.

Symptoms

- feeling dizzy or lightheaded, especially when moving from lying or sitting to standing
- feeling faint or fainting
- blurry or tunnel vision
- feeling nauseated, hot or clammy
- feeling weak and shaky in the legs.

Causes

- · when you stand or sit up suddenly
- in the morning when your blood pressure is naturally lower
- after a large meal or alcohol
- during exercise
- · when you strain on the toilet
- when you're unwell.

Are you at risk?

Some risk factors include:

- high blood pressure
- some medical conditions such as diabetes and heart failure
- some medications such as diuretics, antidepressants and medicines to lower blood pressure
- neurological conditions such as Parkinson's disease and dementia
- dehydration
- anaemia
- excessive alcohol consumption
- · prolonged bed rest or inactivity.



What can you do?

- Speak to your health professional about any symptoms or concerns you have.
- · Discuss the risks and side-effects of your medicines with them.
- Exercise gently before getting up from a chair or bed:
 - move your feet up and down and clench and unclench your hands.
 - get out of bed slowly sit on the side of the bed first.
 - count to 10 slowly, then stand.
- Stand slowly when rising from a chair:
 - make sure you have something to hold on to
 - · wait for a moment before walking.
- Sit down immediately if you feel dizzy.









stayonyourfeet.com.au

For more information or enquiries, contact healthpolicy@health.wa.gov.au





Tinkle Thursday

Urinary incontinence is a risk factor for falls.

Did you know?

One in 14 Australians over the age of 65 and one in 4 over the age of 85 experience severe incontinence – either bowel, urinary or both.

What does it mean?

Urinary incontinence and continence problems may include:

- · stress incontinence
- · urge incontinence
- nocturia (more frequent urination at night)
- overflow incontinence
- function incontinence
- prolapse.

Bowel problems have many causes and may result in:

- faecal incontinence
- constipation
- · diarrhoea.



How to stay healthy

- Talk to your health professional.
- Reduce how much caffeine and alcohol you drink these can upset the bladder.
- Eat plenty of vegetables and fruits to prevent constipation straining can weaken your pelvic floor muscles.
- Rest with your legs up for an hour in the afternoon or evening.
- Have the path to the toilet well-lit at night and free of clutter, so you don't trip.
- Consider using a urinal or commode by the bed at night.
- · Know where the toilets are, and don't rush.

Find out more



continence.org.au



healthywa.wa.gov.au



stayonyourfeet.com.au







Forget Me Not Friday

Cognitive impairment means difficulties in one or more areas of cognition, including memory, thinking, orientation, comprehension, calculation, learning capacity, language and judgement.

Symptoms

- trouble making good decisions
- memory loss affecting daily activities
- problems with abstract thinking
- problems with language (e.g. forgetting words or word replacement)
- loss of initiative or starting things
- changes in personality
- · disorientation to time and place
- · changes in mood or behaviour.

Are you at risk?

Some risk factors include:

- being aged 65 or over
- severe illness
- recent surgery
- history of delirium, dementia or depression
- · hearing or visual impairment
- alcohol or sedative use
- taking several medications.



Know the causes

- being in unfamiliar surroundings
- · during or after an infection
- · experiencing severe pain
- not drinking enough fluids (dehydration) or getting adequate nutrition
- constipation or urine retention
- · taking certain medications, including 'over the counter' medications
- · withdrawal from medication, alcohol or other substances.

What can you do?

- tell your health professional if you experience any changes to your thinking or memory
- if you have a pre-existing condition, such as dementia, sharing your daily routine with others can improve comfort and assist in reducing distress

Family and carers - things to bring

- · well fitting, non-slip shoes or slippers
- any usual glasses or hearing aids (and spare batteries)
- any usual walking aids, as requested
- · nightwear that's above ankle length
- personal items (e.g. photographs, personal clothing).









healthywa.wa.gov.au



